



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Analyzing the Structure and Function of Novel Cytochromes from a Natural Microbial Community

A. Siebers, S. Singer, M. Thelen

August 28, 2006

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

**Analyzing the Structure and Function of Novel Cytochromes from a
Natural Microbial Community**

Anna Siebers, Steve Singer, and Michael P. Thelen

Office of Science, Science Undergraduate Laboratory Internship (SULI)

University of California, San Diego

Lawrence Livermore National Laboratory

Livermore, California

August 30, 2006

UCRL-TR-224023

Prepared in partial fulfillment of the requirements of the Office of Science, Department of Energy's Science Undergraduate Laboratory Internship under the direction of Dr. Michael P. Thelen in the Microbial Biosciences Division at the Lawrence Livermore National Laboratory.

Participant: Anna Siebers

Research Advisor: Michael P. Thelen

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

Table of Contents

Abstract	1
Introduction	2
Materials and Methods	3
Results and Discussion	6
Conclusions	8
Acknowledgements	9
References	10
Figures and Tables	11

ABSTRACT

The Richmond mine in Iron Mountain, California, provides an unusual ecosystem suitable for the growth of microbial biofilms which produce many unique proteins. Through iron oxidation, these proteins facilitate acid mine drainage (AMD). Because this habitat is extremely acidic, survival is an extraordinary feat and the process of environmental selection is rare. In order to understand the mechanisms by which these organisms oxidize iron and gain electrons for energy, biochemical studies were applied. More specifically, column chromatography, spectrophotometry, and gel electrophoresis were used to determine the proteins present in different biofilms. Two specific locations of the mine researched were the AB drift and Ultraback C (UBC), which were both found to contain at least five different types of protein and a large amount of heme-bound cytochromes. Another application of these methods was to investigate proteins playing a major role within the community; one protein selected was cytochrome 579 (Cyt₅₇₉) due to its abundance in the biofilm, iron oxidizing potential, and signature absorbance of 579nm. The structure and function of Cyt₅₇₉ could be characterized by the isolation of its heme, which was completed using column chromatography; however, one of the challenges has been liberating the heme from the column. Further research, including acid-base and temperature profiling of Cyt₅₇₉ should help elucidate its structural changes within alternate environments and metabolism within the community.

INTRODUCTION

Acid mine drainage is a current environmental problem which exists throughout the world. Water flowing through coal or metal mines contributes toxic metals and acidity to downstream aquifers, which in turn pollute water supplies. Biogeochemical studies implicate micro-organisms as a key component in AMD formation. Studies of microbial communities that thrive in these acidic and metal-rich environments may not only lead to remediation strategies which would prevent AMD, but also illuminate the function of these unique, extremophilic organisms. Furthermore, these organisms could potentially be used for industrial purposes, such as in bio-leaching of valuable metals, or in the biotechnological use of highly stable proteins.

Our work focuses on an AMD site in Iron Mountain, California, where a biofilm forms along the surface of the acidic water ($\text{pH} < 1$). This biofilm consists of self-sufficient extremophiles that accelerate the dissolution of the pyrite (FeS_2) bed in the mine. The biofilm gains energy by Fe (II) oxidation and grows autotrophically by fixing CO_2 and N_2 from the air [1]. The acidic mine water also has high concentrations of copper, zinc and arsenic [2].

To understand how the biofilm functions, metagenomic and proteomic studies were conducted. Greater than 12,000 genes within the community were identified by genomic sequencing, and two genomes were then reconstructed for the dominant species, *Leptospirillum* group II and *Ferroplasma* type II [1]. Shotgun proteomics, utilizing two-dimensional mass spectrometry, was used to identify proteins expressed within the biofilm [3]. Proteins from the extracellular and periplasmic regions were studied because exterior proteins not only play a vital role in the metabolism of the organism and interactions within the community, but are also in contact with the extreme environment. The majority of the proteins in the extracellular fraction were hypothetical, meaning they have low sequence similarity to proteins of known function.

Of particular interest was an abundant protein, Cyt₅₇₉, which has a sequence that corresponds to a hypothetical gene of the source organism, *Leptospirillum* group II. Although Cyt₅₇₉ contains a cytochrome c like amino acid sequence motif (CysXaaXaaCysHis), it has little sequence similarity to cytochrome c [4]. This important protein was isolated from environmental biofilm samples and then highly enriched by column chromatography. The visible spectra of Cyt₅₇₉ is indicative of an unusual heme co-factor and reaffirms a cytochrome structure. To study the heme and the specific mechanisms of electron transport from Fe(II) to *Leptospirillum* bacteria, methods were developed to isolate this unusual heme using the model substrate cytochrome c and then the uncharacterized Cyt₅₇₉.

MATERIALS AND METHODS

Heme Extraction from Cytochrome C

To form a solution of free heme and cytochrome c, 48 mg of cytochrome c was first dissolved in 1 ml of water. Then, this was added to a solution of 80 mg of Ag₂So₄ dissolved in 9 ml of water and 0.8 ml of acetic acid. This mixture was incubated in the dark for 4 hours at 40°C and then centrifuged to remove heme aggregates and silver precipitate. See reference [5] for heme extraction method.

The supernatant from the above procedure, containing cytochrome c (protein-bound heme), apo-cytochrome c and free heme, was then separated by SDS-polyacrylamide gel electrophoresis [6] in order to test for efficiency of heme extraction. Four samples were loaded, including two of the previously silver-treated cytochrome c samples and two non-treated, cytochrome c controls (1 mg/mL). One of each was loaded on opposite sides of the gel; the left side was treated with o-dianisidine to stain specifically for heme [7], and the right was stained with Coomassie blue dye for protein.

2 mL of the treated cytochrome c solution was also applied to a SP-sepharose column in order to separate the free heme from the apo-cytochrome c and heme-bound protein and also to collect a sample

of heme. Three buffers were successively passed through the column: 2 M NaCl in 100 mL $\text{NaC}_2\text{H}_3\text{O}_2$ pH 5, 100 mM Na_2CO_3 pH 10, and 0.1 - 0.2 M NaOH.

Furthermore, 1 mL of cytochrome c was also applied to a G-25 Sephadex column in order to obtain a sample of free heme by elution with 0.1 - 0.2 M NaOH.

Acid-Base Profiling of Cytochrome C

For testing structural changes in cytochrome c at different pH values, thirteen buffers were composed (see Table 1) [8]. Each cytochrome c sample was diluted by adding 800 μL of the specific pH buffer to 200 μL of the 1 mg/mL cytochrome c solution. Then, 25 μL of each diluted cytochrome sample was added to 975 μL of pH 7 buffer to make 1 mL samples for spectrophotometry.

Processing the Biofilm: Isolation of Extracellular Fraction

The Ultraback C (UBC) biofilm (collected 6/16/06) was fractioned into extracellular and cellular membrane samples in order to separate the different types of proteins. First, a 50 mL sample of biofilm was thawed by adding a small amount (~ 10 mL) of 0.2 M H_2SO_4 pH 1.1 into the tube and placing it in a beaker of warm water. Once thawed and transferred into a 300 mL glass beaker, it was suspended in 150 mL of 0.2 M H_2SO_4 pH 1.1 and stirred for 10 minutes at room temperature. The solution was homogenized, stirred for 1-1.5 hours at 4 °C, homogenized again, and then centrifuged for 12 minutes at 17500 rpm/rcf and 20 °C in the SS-34 rotor. The supernatant was decanted and the pellet (cellular fraction) was stored in the freezer for further analysis.

25 grams of $(\text{NH}_4)_2\text{SO}_4$ was added per 100 mL of supernatant and this solution was stirred for 1 hour at 4 °C, centrifuged for 15 minutes at 17.5 K rpm/rcf, and decanted (supernatant for 95% fraction). The precipitate from this step made up the 45% precipitate fraction. The pellet was dissolved in 3-5 mL of 20 mM H_2SO_4 /100 mM ammonium sulfate pH 2.2 (buffer A), centrifuged, decanted, and then stored.

Next, 28 grams of $(\text{NH}_4)_2\text{SO}_4$ was added to each 100 mL of supernatant from the 45% precipitate fraction. This mixture was stirred for 2 hours at 4 °C, centrifuged for 20 minutes at 17.5 K rpm/rcf, and decanted. The pellet was dissolved in 3-5 mL of buffer A, which made the 95% precipitate fraction. Then, it was dialyzed overnight in a 500 mL of buffer A. Protein concentrations were estimated by the Bradford protein assay, using bovine serum albumin as a standard [9].

Enrichment of Cyt₅₇₉ and Heme Extraction from Cyt₅₇₉

Cation-exchange chromatography was used in order to enrich Cyt₅₇₉. About 5 mL of the 45% precipitate UBC extracellular fraction was loaded onto an SP-sepharose fast flow column equilibrated with 6 column volumes of buffer A. The absorbed sample was washed with ~3 columns of buffer A and the cytochrome fraction eluted with 100 mM NaOAc pH 5.0. Remaining protein was eluted with a 0-2M NaCl gradient in pH 5.0 buffer.

A G-25 sephadex column was loaded with about 1 mL of treated Cyt₅₇₉ and washed with 100 mL of 100 mM NaOAc pH 5.0 buffer and 100 mL of 1 M NaCl.

RESULTS and DISCUSSION

Heme Extraction Using Cytochrome C as a Model

Cytochrome c is an analog for Cyt₅₇₉ because both have the heme-binding sequence motif, implying that heme is covalently bound through disulfide bonds to the two Cysteines. Because samples of Cyt₅₇₉ were limited, biochemical tests were first performed on cytochrome c. Previous attempts to extract heme from Cyt₅₇₉ were not successful, but a silver sulfate treatment [5] appeared promising. Gel electrophoresis was initially used to verify that the silver sulfate treatment was effective in heme extraction. Following separation of silver-treated and untreated cytochrome c proteins, staining the polyacrylamide gel with Coomassie-blue dye revealed protein in both samples. However, staining the gel with o-dianisidine resulted in the sole appearance of free heme at a position of < 4 kDa, while the non-treated protein appeared at ~16 kDa (**Figure 1**).

The SP-sepharose column eluted apo-cytochrome c protein with the pH 10 buffer, and heme-bound protein and some free heme with 0.1 M NaOH. The majority of free heme remained attached to the column, even after an hour exposure with 0.2M NaOH. The G-25 sephadex column eluted apo-cytochrome c protein immediately after washing pH 5 buffer through the column. Then, 3 heme-bound 3 mL fractions were eluted using 0.1 M NaOH. However, similar to the SP-sepharose column, most of the free heme remained spread out and bound tightly along the column. This is further discussed below in “Heme Extraction from Cyt₅₇₉ and Cytochrome C”.

Acid Base Profiling of Cytochrome C

As an indicator of structure and stability of cytochrome c, visible spectra from 350 to 800 nm were taken after 24 hours incubation in different pH buffers. These spectra indicated that the heme group of cytochrome c was reduced at both pH 9 and 10, whereas no other pH significantly affected the protein. In all cases, the visible spectra indicated the cytochrome remained intact after 24 hours.

Cytochrome c was used again as a model for Cyt₅₇₉, which will soon be tested when more biofilm samples are processed.

Enrichment of Cyt₅₇₉ and Heme Extraction

In the UBC biofilm sample, much more protein from the extracellular fraction was detected in the 45% ammonium sulfate precipitate sample than in the 95% precipitate. In contrast, most of the extracellular protein of previously studied AMD biofilms was recovered from the 95% precipitate fraction. This observation is interesting in that the UBC biofilm compared to previously characterized biofilms perhaps has a different structure, contains different amounts of metals, or that Cyt₅₇₉ carries a different charge. Cation-exchange chromatography of the UBC 45% fraction eluted fractions which were red due to Cyt₅₇₉, yielding 1.5 mg protein from the biofilm sample.

Following silver treatment of Cyt₅₇₉, heme aggregates were seen and the solution turned slightly red probably due to liberated heme (similar to cytochrome c). After gel electrophoresis of the treated sample, heme-bound protein was detected through heme staining (**Figure 2**). However, free heme did not appear to be present. Perhaps this occurred because the heme migrated too far in the gel or the free heme was not concentrated enough for detection. Another gel will be run with Cyt₅₇₉ for a shorter period in order to eliminate one of these possibilities.

The G-25 sephadex column eluted a protein fraction with the pH 5.0 buffer; however, the heme remained bound to the column, even after passing 1M NaCl through the column. Cyt₅₇₉ has not yet been tested in column experiments with NaOH.

Comparison of Cyt₅₇₉ and Cytochrome C Heme Groups

The results from the SP-Sepharose and Sephadex-25 columns demonstrated that the free heme bound tightly to both columns. The dark, red band that remained at the top of these columns could

contain precipitated cytochrome or aggregated heme derived from the silver sulfate treatment. Although the heme structure of Cyt₅₇₉ is probably similar to the common porphyrin ring structure of most heme compounds that are soluble in alkaline solutions, it appears to be unique due to its specific visible (red-shifted) spectral characteristics. One possible explanation of the strong binding of heme to these columns is that it may be structurally altered by the addition of a formyl group at one of its vinyl side chains. This heme alteration theory has been reported for a recombinant cytochrome purified from *E. coli* [10] and confers a higher reduction potential that would make it more efficient in iron oxidation.

CONCLUSIONS

In the process of isolating heme from cytochrome c, the heme-bound protein and some free heme could be recovered using either Sephadex G-25 or SP-sepharose columns with 0.1 M NaOH buffer. Some free heme was also removed from Cyt₅₇₉ using a Sephadex G-25 column, however the yield was low. Other methods for eluting greater quantities of free heme from cytochrome c will be developed prior to isolating the heme from Cyt₅₇₉. Cytochrome c was shown to be stable over a wide range of pH and reduced spontaneously at pH 9 and 10, but not at lower or higher pH values. The potential results of acid-base profiling of Cyt₅₇₉ may exhibit similar or distinct results compared to those of cytochrome c and will allude to the structure and stability of Cyt₅₇₉. Ammonium sulfate treatment of the UBC and AB Drift extracellular fractions resulted in enriched samples of Cyt₅₇₉. Through purification and heme characterization studies, the properties of this unique cytochrome appear to be distinct in different biofilms. This could indicate that the structure of Cyt₅₇₉ is selected for its optimum redox function by the influence of environmental conditions on the microbial community.

ACKNOWLEDGEMENTS

I would like to thank my mentor, Michael P. Thelen, for giving me the opportunity to work in his lab at Lawrence Livermore National Laboratory and also for his enthusiasm, knowledge, and on-going support. Thanks to the awesome Microbial Biosciences crew, including my other mentor whom I worked with daily, Steve Singer, and scientists Mona Hwang and Chris Jeans. They never ceased to answer my frequent questions and created a wonderful learning environment. Thanks to the U.S. Department of Energy, Office of Science for the SULI program and an amazing, educational experience. Lastly, thanks to my favorite high school teacher, Frankie Tate, for encouraging me to apply for the Edward Teller Science Scholarship.

REFERENCES

- [1] Tyson et al (2004) "Community Structure and metabolism through reconstruction of microbial genomes from the environment" *Nature* 428, 37
- [2] J. Banfield and T. Gihring (2006) "Sulfide Mineral Weathering and Acid Mine Drainage Research." <http://seismo.berkeley.edu/~jill/amd/AMDresearch.html>
- [3] Ram et al (2005) "Community Proteomics of a Natural Microbial Biofilm" *Science* 308, 1915-20.
- [4] M.P. Thelen (2006, June 20). *Biofilm Proteins Associated with Acid Mine Drainage*. Presented at American Society of Microbiology in Orlando, Florida.
- [5] W. Fisher, H. Taniuchi and C. Anfinsen (1973) "On the Role of Heme in the Formation of the Structure of Cytochrome c" *Journal of Biological Chemistry* 248, 3188-95.
- [6] U.K. Laemmli (1970) *Nature* 227, 680-5.
- [7] R.T. Francis and R.R. Becker (1984) *Anal. Chemistry* 136, 509-14.
- [8] R. Dawson, D. Elliott, W. Elliott and K. Jones (1986) Data for Biochemical Research New York: Oxford University Press.
- [9] M.M. Bradford (1976) *Anal. Biochemistry* 72, 248-54.
- [10] J. Fisher, T. Todaro, E. Luna, D. Sanders, L. Hunsicker-Wang, K. Patel, K. Bren, E. Gomez-Moran, M. Hill, J. Ai, T. Loehr, W. Oertling, P. Williams, C. Stout, D. McRee and A. Pastuszyn (2004) "Cytochrome rC₅₅₂, Formed during Expression of the Truncated, *Thermus thermophilus* Cytochrome c₅₅₂ Gene in the Cytoplasm of *Escherichia coli*, Reacts Spontaneously To Form Protein-Bound 2-Formyl-4-vinyl (*Spirographis*) Heme" *Biochemistry* 43, 12162-76.
- [11] S. Singer (2006, July 26). *Understanding Natural Microbial Communities Using Proteogenomics and Biochemistry*. Presented at the Annual Biosciences Directorate Postdoctoral Symposium, Lawrence Livermore National Laboratory.
- [12] J. Banfield, N. Verberkmoes, R. Hettich, M. Thelen (2005) *OMICS A Journal of Integrative Biology* 9, 301-33.
- [13] B. Baker and J. Banfield (2002) *FEMS Microbiology Ecology* 44, 139-52.




Figure 1. Heme stain of pure Cytochrome C




Figure 2. Protein stain (1) and heme stain (2) of enriched Cyt₅₇₉

Table 1- Buffers

Buffer Type	Components
pH 1 KCl/HCl	2.5 mL 0.2 M KCl 6.7 mL 0.2 M HCl 0.8 mL H ₂ O
pH 2 KCl/HCl	2.5 mL 0.2 M KCl 6.5 mL 0.2 M HCl 1.0 mL H ₂ O
pH 3 Glycine-HCl	2.5 mL 0.2 M glycine 0.57 mL 0.2 M HCl 6.93 mL H ₂ O
pH 4 NaC ₂ H ₃ O ₂	1.8 mL 0.2 M NaOAc 8.2 mL 0.2 M HOAc
pH 5 NaC ₂ H ₃ O ₂	7.0 mL 0.2 M NaOAc 3.0 mL 0.2 M HOAc
pH 6 MES-NaOH	2.5 mL 0.1 M MES 0.8 mL 0.1 M NaOH 1.7 mL H ₂ O
pH 7 KH ₂ PO ₄ -NaOH	5 mL 0.1 M KH ₂ PO ₄ 2.91 mL 0.1 M NaOH 2.09 mL H ₂ O
pH 8 KH ₂ PO ₄ -NaOH	5 mL 0.1 M KH ₂ PO ₄ 4.61 mL 0.1 M NaOH 0.39 mL H ₂ O
pH 9 Glycine-NaOH	2.5 mL 0.2 M glycine 0.44 mL 0.2 M NaOH 7.06 mL H ₂ O
pH 10 Glycine-NaOH	2.5 mL 0.2 M glycine 1.6 mL 0.2 M NaOH 5.9 mL H ₂ O
pH 11 Phosphate	5 mL 0.005 Na ₂ HPO ₄ 0.41 mL 0.1 M NaOH 4.59 mL H ₂ O
pH 12 Hydroxide-chloride	2.5 mL 0.2 M KCl 0.6 mL 0.2 M NaOH 6.9 mL H ₂ O
pH 13 Hydroxide-chloride	2.5 mL 0.2 M KCl 6.6 mL 0.2 M NaOH 0.9 mL H ₂ O

